

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## NOTES ON FLORIDA FUNG!.--14.

BY W. W. CALKINS, CHICAGO, ILLINOIS.

- 251. Polyporus nivosus, B. & Br.—Found on dead gum trees along with P. hemileucus; rare.
  - 252. POLYPORUS LINDBLADII, B. var. Very rare on old pine logs.
- 253. POLYPORUS FERRUGINOSUS, Fr.—Rare and only found once on an old log. Looks quite different from P. obliquus. The above three species have been passed upon by Dr. Cooke.
  - 254. Kneiffia ambigua, Karsten.—On an oak log; rare.
  - 255. DIATRYPELLA DEUSTA, E. & M.—Abundant on palmetto stems.
  - 256. DIATRYPELLA VERRUCÆFORMIS, Ehr.—On dead limbs.
  - 257. DIATRYPELLA OPACA, Cke -On dead holly.
  - 258. Coniosporium Arundinis, Cda.—On Sabal stems.
  - 259. UREDO FICUS, Cast.—Abundant on fig leaves.
- 260. OIDIUM MEGALOSPORUM, B.—Abundant on rotten logs in swamps.
  - 261. SEPTORIA HYDROCOTYLES, E. & M.—Abundant.
  - 262. Septoria Symploci, E. & M.—On leaves of the sugar tree.
  - 263.
  - SEPTORIA SERPENTARIA, E. & M.—On Olea leaves. SEPTORIA NIPHOSTOMA, B. & C.—On Magnolia leaves. TRABUTIA QUERCINA, Fr. & R.—Abundant. MBLANCONIUM SABAL, Cke.—On Sabal stems. 264.
  - 265.
  - 266. 267.
  - Puccinia Hydrocotyles, Mont.—Abundant. 268. MACROSPORIUM NERII, Cke.—Abundant on fallen leaves of
- Oleander. 269. Fusarium Yuccæ, Cke.—On Yucca aloifolia leaves.
- 270. HELICOTRICHUM OBSCURUM, Cda.—Abundant on fallen Persea leaves; very obscure, too.
  271. TREMELLA MESENTERICA, Fr.—Occasional on dead limbs.

  - 272. TREMELLA FOLIACEA, Pers.—Abundant on rotten limbs.
  - 273.
  - PHYLLACHORA DEMERSA, Cda.—On Persea leaves.
    PHOMA LEGUMINUM, West.—On pods of Glottidium. 274.
  - PHOMA CLITORIACARPA. Cke.—On pods of Glottidium. MBLOGRAMMA FULIGINOSUM, Ell.—On Sabal stems. **275.**
  - 276.

  - 277. Russula emetica. Fr.—Abundant in woods. 278. Endothia gyrosa, Sw.—On old oak limbs; abundant.
- 279. Geaster hygrometricus, Pers.-Very common in sandy fields. 280. Anthostomella minor, E & M.—A new species. Journ. Mycol. Vol. III, p. 43. Common on Sabal stems.

## NEW LITERATURE.

BY W. A. KELLERMAN.

"GERMINATION OF ERGOT FROM THE WILD RYE." B. D. Halsted, in Bulletin of the Iowa Agricultural College, Botanical Dept. November, 1886.

Ergotted grains from Elymus Canadensis were placed in a large flower pot on March 10th. Two months later, growth became evident

and in a week the heads were formed and became stalked. On May 31st, the grains had mature heads and were then examined microscopically. The stems were two to three centimeters long and the ascigerous heads two to three millimeters in diameter and sphæroidal. The asci were 4 x 230 u and 8-spored; the spores were 1 x 120 u. Saccardo gives the size of spores of Claviceps purpurea (Fr.) as 50-76 u. Dr. Halsted finds that the ascigerous stems from the cultivated rye as figured by Dr. Winter (Pilze II, 91) are much shorter and the heads larger than those mentioned above. That the species of ergot infecting the wild rye is the same as that of the cultivated rye, Claviceps purpurea, remains an open question.

"Notes upon the Peronosporeæ for 1886." l. c.

"Notes upon the Ustilagineæ." 1. c.

"THE ASH-LEAF RUST. ' .1. C.

"THE CLOVER MOULD (Cladosporium). l. c.

"Fungi of Forest Trees." 1. c.

"COLORADO FUNGI." July, 1886. 1. c.

Twenty-five species are enumerated, of these, the following new: Tubercularia Lupini, Farlow, in litt, on L. Kingii, Watson, Gunnison, Colo.; Puccinia sp., on Artemisia Mexicana, Willd., Aecidium and Puccin-The æcidia are long and slender and perhaps new. Of the Puccinia, Dr. Farlow says; "It is not the usual form on Artemisa and does not exactly correspond to anything I have examined. \* \* I think it may be new and the æcidium may very likely be its æcidium." A parasitic fungus was found on Pseudotsuga Louglasii, Carr, causing distortions called "eagle nests." Of this, Dr. Farlow says: "It has almost a greater similarity to some of the Cæomata and Abies than to a Peridermium; but it has a peridium more delicate than in any species known. It seems to be new. In habit and the distinctions produced, it reminds one of Aecidium corruscens, Fr., but unless it changes very much as it grows older, it cannot be that species. It is characterized by the size of its spores and size and thin markings of peridial cells."

"RELATION BETWEEN 'CEDAR-APPLES' AND THE LEAF RUST ON THE WILD CRAB APPLE." 1 c.

PARTIAL LIST OF IOWA POWDERY MILDEWS." A. S. Hitchcock.

1. c.

"Fungi injurious to Grasses and Clovers." Wm. Trelease, St. Louis, Mo. From "Beal's Grasses of North America."

A popular account, with illustrations of fourteen species, covering pp. 413-421.

"UEBER EINIGE AUF RUBUS ARCTICUS L. VORKOMMENDE PARASITISCHE PILZE." Af G.Lagerheim. Separataftryck ur Botaniska Notisier, 1887. "ESPERIMENTI SUL PARASITISMO DELL' AGARICUS MELLEUS, VAHL.

Di L. Savastano. Nuovo Giornale Batanico Italiano April, 1887. "Botanical Manuals for Students." Chas. E. Bessey. American Naturalist, April, 1887.

"St. George's Mushroom, Agaricus gambosus," Fr. Worthington G. Smith, Gardner's Chronicle, April 23, 1887.
"An Introduction to the Study of Lichens." By Henry Willey.

Pp. 72, ten plates.